

MEETING ABSTRACT

Open Access

Can focused echocardiographic evaluation in life support (FEEL) be used to predict resuscitation outcome or termination of resuscitation (TOR)? A prospective trial

JY Suh*, YS Jo, JH Choi, HM Kim

From 10th WINFOCUS World Congress on Ultrasound in Emergency and Critical Care Kuala Lumpur, Malaysia. 16-19 November 2014

Background

There are many studies about predictive factor of return of spontaneous circulation (ROSC) during the cardiopulmonary resuscitation (CPR), but no definite guideline or predictive factor for termination of CPR.

Objective

The purpose of this study is to figure out the relationship between ROSC and cardiac activity findings by echocardiography(echo) in cardiac arrest patients, and investigate the cardiac standstill period can be used as indicator for termination resuscitation.

Patients and methods

We performed a prospective, observational study of non-consecutive, non-trauma, adult, out of hospital cardiac arrest (OHCA) patients. This study conducted in emergency department of single tertiary university hospital. Echo performed every 2 minute simultaneously with pulse check within 10 second throughout the arrest, which was managed by the usual advanced cardiac life-support treatment guidelines. We obtained sub-xiphoid or parasternal long axis view of echo. We defined echocardiographic evidence of cardiac kinetic activity as any detected motion of the myocardium, ranging from visible ventricular fibrillation to coordinated ventricular contractions.

Results

During 10 months, 33 patients were enrolled in the study. Witness arrest patients were 22 (66.7%), 12 (36.3%)

patients received bystander CPR. Interval from EMS call to EMS scene arrival is 8 ± 3 minutes, Interval from EMS call to EMS hospital arrival is 23 ± 6 minutes. 17 (51.5%) patients attained ROSC, and 1 (3%) patient discharged in cerebral performance category 1. Repetitive cardiac standstill periods in ROSC group and no ROSC group were 2.9 ± 2.4 minutes versus 18.9 ± 8.4 minutes, respectively ($p<0.001$). Cardiac standstill period over 8 minutes




Figure 1

* Correspondence: hongwe@naver.com

Emergency medicine, Soonchunhyung University Bucheon Hospital, Bucheon, South Korea

predicted eventual no ROSC with a sensitivity of 86.7% and a specificity of 100%. A receiver operating curve was generated to determine the accuracy of echocardiographic cardiac standstill period for predicting no ROSC. The Area under the ROC curve was 0.985 ($p < 0.01$). Figure 1.

Conclusion

If over 8 minutes repetitive cardiac standstill was identified by echo during the CPR, termination of resuscitation could be considered.

Published: 9 March 2015

doi:10.1186/2036-7902-7-S1-A24

Cite this article as: Suh *et al.*: Can focused echocardiographic evaluation in life support (FEEL) be used to predict resuscitation outcome or termination of resuscitation (TOR)? A prospective trial. *Critical Ultrasound Journal* 2015 **7**(Suppl 1):A24.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
